

## 参考文献

- [1] Peek RM, Crabtree JE. Helicobacter infection and gastric neoplasia [J]. The Journal of pathology, 2006, 2008(2): 233-248.
- [2] Wang M, Chen C, Gao X, et al. Distribution of *Helicobacter pylori* virulence markers in patients with gastroduodenal diseases in a region at high risk of gastric cancer [J]. Microbial pathogenesis, 2013, 59: 13-18.
- [3] Salimzadeh L, Bagheri N, Zamanzad B, et al. Frequency of virulence factors in *Helicobacter pylori*-infected patients with gastritis [J]. Microbial pathogenesis, 2015, 80: 67-72.
- [4] Umit H, Tezel A, Bukavaz S, et al. The relationship between virulence factors of *Helicobacter pylori* and severity of gastritis in infected patients [J]. Digestive Diseases and Sciences, 2009, 54(1): 103-110.
- [5] Boughan PK, Argent RH, Body-Malapel M, et al. Nucleotide-binding Oligomerization Domain-1 and Epidermal Growth Factor Receptor Critical Regulators of  $\beta$ -Defensins during *Helicobacter pylori* infection [J]. Journal of Biological Chemistry, 2006, 281(17): 11637-11648.
- [6] Yamaoka Y, Kodama T, Kita M, et al. Relationship of *vacA* genotypes of *Helicobacter pylori* to *cagA* status, cytotoxin production, and clinical outcome [J]. Helicobacter, 1998, 3(4): 241-253.
- [7] Park H, Li Z, Yang X O, et al. A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17 [J]. Nature immunology, 2005, 6(11): 1133-1141.
- [8] Ivanov II, McKenzie BS, Zhou L, et al. The orphan nuclear receptor ROR  $\gamma$  t directs the differentiation program of proinflammatory IL-17<sup>+</sup> T helper cells [J]. Cell, 2006, 126(6): 1121-1133.
- [9] Ouyang W, Kolls JK, Zheng Y. The biological functions of T helper 17 cell effector cytokines in inflammation [J]. Immunity, 2008, 28(4): 454-467.
- [10] Liang SC, Long AJ, Bennett F, et al. An IL-17F/A heterodimer protein is produced by mouse Th17 cells and induces airway neutrophil recruitment [J]. The Journal of Immunology, 2007, 179(11): 7791-7799.
- [11] Iwakura Y, Ishigame H. The IL-23/IL-17 axis in inflammation [J]. Journal of Clinical Investigation, 2006, 116(5): 1218-1222.
- [12] Kandulski A, Wex T, Kuester D, et al. Naturally occurring regulatory T cells (CD4<sup>+</sup>, CD25<sup>high</sup>, FOXP3<sup>+</sup>) in the antrum and cardia are associated with higher *H. pylori* Colonization and Increased Gene Expression of TGF- $\beta$ . [J]. Helicobacter, 2008, 13(4): 295-303.
- [13] Gil JH, Seo JW, Cho MS, et al. Role of Treg and TH17 cells of the gastric mucosa in children with *Helicobacter pylori* gastritis [J]. Journal of pediatric gastroenterology and nutrition, 2014, 58(2): 252-258.
- [14] Bagheri N, Azadegan-Dehkordi F, Shirzad H, et al. The biological functions of IL-17 in different clinical expressions of *Helicobacter pylori*-infection [J]. Microbial pathogenesis, 2015, 81: 33-38.
- [15] Kao JY, Zhang M, Miller MJ, et al. *Helicobacter pylori* immune escape is mediated by dendritic cell-induced Treg skewing and Th17 suppression in mice [J]. Gastroenterology, 2010, 138(3): 1046-1054.
- [16] Serrano C, Wright SW, Bimeczok D, et al. Downregulated Th17 responses are associated with reduced gastritis in *Helicobacter pylori*-infected children [J]. Mucosal immunology, 2013, 6(5): 950-959.
- [17] Bagheri N, Azadegan-Dehkordi F, Sanei H, et al. Associations of a TLR4 single-nucleotide polymorphism with *H. pylori* associated gastric diseases in iranian patients [J]. Clinics and research in hepatology and gastroenterology, 2014, 38(3): 366-371.
- [18] Fox JG, Wang TC. Inflammation, atrophy, and gastric cancer [J]. Journal of Clinical Investigation, 2007, 117(1): 60-69.
- [19] Shi Y, Liu XF, Zhuang Y, et al. *Helicobacter pylori*-induced Th17 responses modulate Th1 cell responses, benefit bacterial growth, and contribute to pathology in mice [J]. The Journal of Immunology, 2010, 184(9): 5121-5129.
- [20] Rahimian G, Sanei MH, Shirzad H, et al. Virulence factors of *Helicobacter pylori vacA* increase markedly gastric mucosal TGF- $\beta$ , mRNA expression in gastritis patients [J]. Microbial pathogenesis, 2014, 67: 1-7.
- [21] Mizuno T, Ando T, Nobata K, et al. Interleukin-17 levels in *Helicobacter pylori*-infected gastric mucosa and pathologic sequelae of colonization [J]. World Journal of Gastroenterology, 2005, 11(40): 6305-6311.
- [22] Sebkova L, Pellicanò A, Monteleone G, et al. Extracellular signal-regulated protein kinase mediates interleukin 17 (IL-17)-induced IL-8 secretion in *Helicobacter pylori*-infected human gastric epithelial cells [J]. Infection and Immunity, 2004, 72(9): 5019-5026.
- [23] Siomek A, Rytarowska A, Szaflarska-Poplawska A, et al. *Helicobacter pylori* infection is associated with oxidatively damaged DNA in human leukocytes and decreased level of urinary 8-oxo-7, 8-dihydroguanine [J]. Carcinogenesis, 2006, 27(3): 405-408.
- [24] Amedei A, Cappon A, Codolo G, et al. The neutrophil-activating protein of *Helicobacter pylori* promotes Th1 immune responses [J]. Journal of Clinical Investigation, 2006, 116(4): 1092-1101.
- [25] Hida N, Shimoyama T, Neville P, et al. Increased expression of IL-10 and IL-12 (p40) mRNA in *Helicobacter pylori* infected gastric mucosa: relation to bacterial *cag* status and peptic ulceration [J]. Journal of clinical pathology, 1999, 52(9): 658-664.
- [26] Figura N, Valassina M, Roviello F, et al. *Helicobacter pylori cagA* and *vacA* types and gastric carcinoma [J]. Digestive and Liver Disease, 2000, 32: 182-183.
- [27] Gil JH, Seo JW, Cho M S, et al. Role of Treg and TH17 cells of the gastric mucosa in children with *Helicobacter pylori* gastritis [J]. Journal of pediatric gastroenterology and nutrition, 2014, 58(2): 245-251.
- [28] Li Q, Li Q, Chen J, et al. Prevalence of Th17 and Treg cells in gastric cancer patients and its correlation with clinical parameters [J]. Oncology Reports, 2013, 30(3): 1215-1222.
- [29] Wu X, Zeng Z, Xu L, et al. Increased expression of IL17A in human gastric cancer and its potential roles in gastric carcinogenesis [J]. Tumor Biology, 2014, 35(6): 5347-5356.
- [30] Pinchuk IV, Morris KT, Nofchissey RA, et al. Stromal cells induce Th17 during *Helicobacter pylori* infection and in the gastric tumor microenvironment [J]. PLoS One, 2013, 8(1): e53798.
- [31] Shibata T, Tahara T, Hirata I, et al. Genetic polymorphism of interleukin-17A and-17F genes in gastric carcinogenesis [J]. Human Immunology, 2009, 70(7): 547-551.
- [32] Qinghai Z, Yanying W, Yunfang C, et al. Effect of interleukin-17A and interleukin-17F gene polymorphisms on the risk of gastric cancer in a Chinese population [J]. Gene, 2014, 537(2): 328-332.
- [33] Serelli-Lee V, Ling KL, Ho C, et al. Persistent *Helicobacter pylori* specific Th17 responses in patients with past *H. pylori* infection are associated with elevated gastric mucosal IL-1 $\beta$  [J]. PLoS One, 2012, 7(6): e39199.
- [34] Lundgren A, Suri-Payer E, Enarsson K, et al. *Helicobacter pylori*-specific CD4<sup>+</sup> CD25<sup>high</sup> regulatory T cells suppress memory T-cell responses to *H. pylori* in infected individuals [J]. Infection and immunity, 2003, 71(4): 1755-1762.
- [35] Harris PR, Wright SW, Serrano C, et al. *Helicobacter pylori* gastritis in children is associated with a regulatory T-cell response [J]. Gastroenterology, 2008, 134(2): 491-499.

(收稿日期:2015-04-15)